

ANAPHYLAXIS MANAGEMENT: EPI & LORATADINE PROTOCOLS

EPINEPHRINE NON AUTO-INJECTOR ADMINISTRATION PROCEDURE

When the listed Indications are met, Contraindications are ruled out and Conditions achieved, in accordance with PEAK's Anaphylaxis Management: Epinephrine & Loratadine Protocols Pre-Administration Checklist (see reverse), a Qualified Provider¹ may administer epinephrine intramuscularly (IM) or subcutaneously (SC) to patients suspected of experiencing an anaphylactic reaction.

STEP 1	STEP 2	STEP 3	STEP 4	STEP 5
VERIFY NEED	PREP INJECTION SITE	VERIFY DRUG	VERIFY DOSAGE	ADMINISTER
<ul style="list-style-type: none"> ✓ Verify need for epinephrine as per the Anaphylaxis Management: EPI & Loratadine Protocols Pre-Administration Checklist (see reverse) 	<ul style="list-style-type: none"> ✓ Upper, lateral thigh <li style="text-align: center;">OR ✓ Upper, lateral arm ✓ Cleanse skin with alcohol prep 	<ul style="list-style-type: none"> ✓ 1:1000 epinephrine ✓ Check expiration ✓ Check ampoule volume clarity 	<ul style="list-style-type: none"> ✓ Draw up the appropriate dose of epinephrine utilizing only a PEAK Epinephrine Syringe <p>CHILD (C) 15 to 30 kg/66 lbs.</p> <ul style="list-style-type: none"> • 0.15 ml (C) <p>ADULT (A) ≥ 30 kg/66 lbs.</p> <ul style="list-style-type: none"> • 0.3 ml (A) 	<ul style="list-style-type: none"> ✓ Insert needle at a 90° degree angle ✓ Slowly push the plunger to administer the medication ✓ Fully depress the plunger to activate automatic needle retraction while needle is still in the patient

¹ Qualified Provider - an individual who has received specific PEAK training