



EMERGENCY RESPONSE TRAINING
A division of Peak Project Management, Inc.

SUSPECTED ANTERIOR SHOULDER DISLOCATION CRITICAL INTERVENTION

PRE-INTERVENTION CHECKLIST

INCIDENT INFORMATION			
PATIENT'S NAME	CHECKLIST COMPLETED BY (PROVIDER)	INCIDENT DATE (DD / MM / YY)	TIME (24 hr)

When the listed Indications for Use are met, Contraindications of Use are ruled out and Conditions of Use achieved, an NUEC 3 certified Provider may perform, with the patient's consent, the 'Suspected Anterior Shoulder Dislocation Critical Intervention'.

INDICATIONS FOR USE

1. Suspected anterior shoulder dislocation in the non-urban environment where transport to definitive medical care is greater than 1 hour from the time of injury; **AND**
2. Inability to safely transport a patient in the position found; **AND/OR**
3. The patient's extremity distal to the injury is exhibiting signs of neurovascular compromise

CONTRAINDICATIONS OF USE

1. Suspected fracture
2. The patient does not consent to the intervention

CONDITIONS OF USE

- Completed a Primary Assessment
- Provided and maintained all necessary critical interventions
- Activated the local emergency response system and notified EMS
- Conducted a Secondary Assessment
- Obtained and recorded baseline vital signs including: GCS, RR, HR, Skin Temp. and Condition
- Ruled out contraindications
- The intervention is to be carried out in accordance with PEAK's 'Suspected Anterior Shoulder Dislocation Critical Intervention Procedure'